arXiv:cond-mat/0401151v1 [cond-mat.mes-hall] 9 Jan 2004

Spin-Orbit Scattering and Time-Reversal Symmetry: Detection of a Spin by Tunneling

M. B. Hastings
Center for Nonlinear Studies and Theoretical Division, Los Alamos National Laboratory, Los Alamos, NM 87545,
hastings@cnls.lanl.gov

(January 6, 2004)

We consider the possibility of detecting spin precession in a magnetic field by nonequilibrium transport processes. We find that time reversal symmetry imposes strong constraints on the problem. Suppose the tunneling occurs directly between systems at two different chemical potentials, rather than sequentially via a third system at an intermediate chemical potential. Then, unless the magnetic fields are extremely strong or spin polarized electrons are used, the periodic signal in the current results from beating together two different precession frequencies, so that observing a signal near the Larmor frequency in this case requires having some cluster with a g factor close to zero.

PACS Numbers: 03.65.Xp,73.23.-b,73.63.Kv

Transport of spin-polarized current through an STM [1,2] opens the possibility of investigating magnetic systems at a much smaller scale than is possible using electron spin resonance [3]. The ability to manipulate single spins via STM is a key part of the development of spintronics [4] and quantum information processing.

Interestingly, even using spin-unpolarized electrons, a periodic modulation of tunneling current through an STM tip was observed experimentally [5]. The original proposed explanation depended on periodic modulation of charge at the Larmor frequency by a precessing spin center. Another very interesting possibility depends on spin-orbit scattering [6–8]. However, in this paper, we will show that, with two exceptions, time reversal symmetry imposes strong constraints on the problem, and for both of the mechanisms considered leads to modulation at a frequency resulting from beating together two different precession frequencies. Then, for a single impurity spin, the modulation in the current appears at twice the Larmor frequency, and not at the Larmor frequency itself. To obtain a signal at the Larmor frequency requires two spins, one of which has a vanishing g-factor [9].

The first possible exception is to have a sufficiently large magnetic field to introduce Aharonov-Bohm phases for the electrons; in this case time-reversal symmetry of the tunneling Hamiltonian is broken by the external field. This case is unlikely to be experimentally relevant, due to the strength of fields required. The more interesting exception involves sequential tunneling through a quantum dot [6,10], and will be discussed more below. Sequential tunneling refers to a system in which there are two leads and a quantum dot, with the chemical potential one lead far above that of on the dot, and the chemical potential on the other lead far below that of the dot. Thus, this system has three different chemical potentials; the results in this paper will apply to systems in which there are only two different chemical potentials.

As a model, we first consider a tunnel junction with a single nearby spin. The electrons can couple to the spin via exchange coupling, and there may also be a spin-orbit coupling present. This provides a means of measuring the spin in terms of its effect on the tunnel current. We first consider the appropriate Hamiltonian for the system in the absence of any external magnetic field, so that the system is time reversal invariant, and then consider the effect of any applied magnetic field. We find that the effect of time reversal symmetry strongly limits the allowed modulation of the current by the spin. Our physical setup with an impurity spin and a single tunnel junction will be similar to that considered in [13]; similar results with some exceptions hold for tunneling through a quantum dot as considered in [6], as discussed below.

Time Reversal Symmetry— Consider the most general possible Hamiltonian, written as $\mathcal{H} = \mathcal{H}_L + \mathcal{H}_R + \mathcal{H}_T$, where $\mathcal{H}_{L,R}$ are the Hamiltonians for the left and right leads of the system and $\mathcal{H}_T$ includes the tunneling terms, as well as terms coupling the electrons to the spin. The two leads may be taken to be Luttinger liquids. There is a voltage difference V between the leads, giving a chemical potential difference $\mu = qV$, where q is the electron charge. We assume that the effects of electron interaction are negligible in the tunneling region, as the tunneling process is weak and only one electron tunnels at a time. Thus, we restrict to terms which are at most second order in the Fermion operators in $\mathcal{H}_T$.

Then, given the requirement of time reversal symmetry, the most general possible $\mathcal{H}_T$, is

$$\mathcal{H}_T = t_0 T_0 + t_1 J_0^j v_j + t_2 J_0 (S_j w_j) + J_{\beta\gamma} (\psi_\beta^\mu)^\dagger S^{\mu\nu} \psi_\gamma^\nu, \quad (1)$$

where a sum over repeated indices is assumed. Here S_j (for $j = x, y, z$) is the spin operator on the impurity spin (which is taken to have total spin S, where S may be larger than $1/2$) and $S^{\mu\nu} = \sigma_j^{\mu\nu} S_j$, where σ_j are the Pauli spin matrices. The letters β, γ identify the particular lead: $\beta, \gamma = L, R$, while ψ^μ are electron destruction operators with spin $\mu = \uparrow, \downarrow$. We have defined the following operators: first, a tunneling opera-

tor $T_0 \equiv (\psi_L^\mu)^\dagger \psi_R^\mu + h.c.$ Next, an electron spin current $J_0^j \equiv i\sigma_j^{\mu\nu}[(\psi_L^\mu)^\dagger \psi_R^\nu - h.c.]$. Finally, an operator related to the current operator, $J_0 \equiv i[(\psi_L^\mu)^\dagger \psi_R^\mu - h.c.]$. The term in t_0 is a tunneling term. The term in t_1, which combines spin-flip with tunneling, can arise via a mechanism similar to that generating spin scattering by nonmagnetic impurities in semiconductors [12], as noted in [6]. The fixed vector v_j is determined by the spin-orbit scattering. The 4 couplings $J_{\beta\gamma}$ are exchange couplings, while the fixed vector w_j is set by both spin-orbit and exchange interactions.

The terms in (1) are time reversal symmetric [11]. This operation complex conjugates all terms in the Hamiltonian, and sends $\psi^\uparrow \to \psi^\downarrow$ and $\psi^\downarrow \to -\psi^\uparrow$, and similarly for $\psi^\dagger$. Further, it changes the sign of the spin operator, $S_i \to -S_i$. Then, J_0 changes sign under time reversal as does S, while J_0^j is time reversal invariant.

The coupling of the tunneling current to the spin depends on a combination of exchange interaction between the electrons and spin with spin-orbit scattering. Since both of these interactions are in the term with t_2, we focus on the following Hamiltonian:

$$\mathcal{H}_e = \mathcal{H}_L + \mathcal{H}_R + t_0 T_0^{\mu\mu} + t_2 J_0 S_z \tag{2}$$

Here, we have chosen w to lie along the z-direction. However, many of the results we obtain will be valid for much more general Hamiltonians, as will be discussed at the end.

External Magnetic Fields— To Eq. (2), we add an external magnetic field. This field breaks the time reversal symmetry and allows for terms which are odd under time reversal, so that $\mathcal{H} = \mathcal{H}_e + \mathcal{H}_o$. Two particularly important allowed terms are

$$\mathcal{H}_o = B_i S_i + t_3 J_0. \tag{3}$$

The first term is the coupling of the field to the spin, while the second results from a possible Aharonov-Bohm phase on the electrons traversing the tunnel junction: the phase of the term coupling to S_z has been changed by the magnetic field, so that it is no longer imaginary. We expect that terms in t_3 can be disregarded for experimentally relevant fields.

The possible introduction of terms like t_3 is very important. To see this importance, return to the Hamiltonian $\mathcal{H}_e$. In this Hamiltonian, the only spin operator that appears is the operator S_z. Thus, if we pick a basis for S that diagonalizes S_z we can consider S to be constant in time, and thus S_z in the Hamiltonian (2) can be considered as a classical parameter. Then, this Hamiltonian is equal to $\mathcal{H}_L + \mathcal{H}_R + [z(\psi_L^\mu)^\dagger \psi_R^\mu + h.c.]$, where $z = t_0 + it_2 S_z$. Since the phase of z is unimportant, physical parameters like the current can depend only on the modulus: $|z| = |t_0 + it_2 S_z| = t_0^2 + t_2^2 S_z^2$.

Thus, the current depends on S_z^2, but cannot depend on the sign of S_z. In fact, for the most general Hamiltonian of Eq. (1), this remains true, as will be discussed more below. On the contrary, if we consider the combined Hamiltonian, $\mathcal{H}_e + t_3 J_0$, the modulus becomes $t_0^2 + t_3^2 + 2t_2 t_3 S_z + t_2^2 S_z^2$, and thus does depend on the sign of S_z. This difference in the dependence of the current on S_z, when S_z is considered as a classical field (unchanging in time) determines, as we will see, how strong the modulation of the current is at the Larmor frequency, ω_L, when S_z acquires time dynamics due to the introduction of an external magnetic field. The Hamiltonian $\mathcal{H}_e + \mathcal{H}_o$, with $t_3 \neq 0$, has been considered previously in the context of a two level system rather than a spin, in which case such terms do not violate time reversal symmetry [13], as well as in the context of a spin system for which such terms do violate time reversal symmetry [7,8]. These authors found a modulation in the tunneling current at the Larmor frequency. In the case of tunneling through a dot, as considered in [6], similar time reversal symmetry violating terms were considered in the matrix $\Omega_{rss'}$ of that paper, again finding modulation at the Larmor frequency.

Current Modulation at Weak Field— We now consider the case instead with a sufficiently weak field, so that t_3 can be disregarded: $\mathcal{H} = \mathcal{H}_e + B_i S_i$. We will find only a very weak modulation of the current at the Larmor frequency, and a much more substantial modulation at twice the Larmor frequency. This has important experimental implications for the detection of a precessing spin with a tunneling current: for small t_3, a much stronger signal should be found at twice the Larmor frequency in this setup with a single tunnel junction.

The effective dynamics of the density matrix for the spin can be found using previous results [15], based on a Keldysh technique [16]. Throughout, we work perturbatively in the tunneling, $\mathcal{H}_T$. Define $I(\mu) = 2\pi\rho_L\rho_R\mu^{\alpha+1}/\Gamma(\alpha+2)$. The parameter α depends on the Luttinger parameters of the leads and the sample geometry; it is equal to zero for Fermi liquid leads [14]. The product of the density of states in the left and right leads at energy μ is $\rho_L\rho_R\mu^\alpha$. Then, the result for the effective dynamics is

$$\dot{\rho} = -i\Big[\mathcal{H}_0, \rho\Big] + \frac{t_2}{2}\frac{\mathrm{d}I}{\mathrm{d}\mu}\Big[S_z, \Big\{\Lambda, \rho\Big\}\Big] - t_2^2\frac{I(\mu)}{2}\Big[S_z, \Big[S_z, \rho\Big]\Big], \tag{4}$$

where $\Lambda = [\mathcal{H}_0, t_2 S_z]$, and $\mathcal{H}_0$ is the Hamiltonian for the spin. This includes the external field, BS_x, as well as an additional effective field along the z direction due to the coupling to the electron current. We choose to absorb the additional field into a renormalization of the bare field, so that the net field is along the x-direction: $\mathcal{H}_0 = BS_x$. Thus, $\Lambda = -it_2 B S_y$.

The density matrix evolution (4) describes the Hamiltonian evolution of the spin under $\mathcal{H}_0$, as well as dissipation (the second term) and decoherence (the third term). For large V, Eq. (4) leads to an effective temperature $T_{\text{eff}} = (\alpha+1)^{-1} qV/2$. This gives a stationary distribu-

tion of the density matrix for the spin:

$$\rho = \frac{1}{2S+1}(1 - BS_x/T_{\rm eff}). \quad (5)$$

Current— We now consider the average current across the system. The current operator J is defined by taking a derivative of the Hamiltonian with respect to an external gauge field, giving

$$J = q(t_0 J_0 - t_2 T_0 S_z). \quad (6)$$

The average current is, for $B << \mu$, [15]

$$\langle J\rangle = q\langle t_0^2 + t_2^2 S_z^2\rangle I(\mu) - qBt_2^2\langle [S_z, S_x]S_z\rangle \frac{{\rm d}I(\mu)}{{\rm d}\mu}. \quad (7)$$

The first term is the current that results from considering S_z to be a classical object, while the second term is a weak quantum correction that slightly reduces the current due to fluctuations in S_z.

As long as the net magnetic field is along the x-direction, $\langle S_z\rangle$=0, and $\langle S_z^2\rangle = (1/3)S(S+1)$. If the field acquires a z-axis component, then, for $S > 1/2$, the expectation value $\langle S_z^2\rangle$ will increase, and thus the net current will increase. The expectation value $\langle [S_z, S_x]S_z\rangle = (B/T_{\rm eff})(1/6)S(S+1)$. Thus, for a net field along the x-direction, the net current is

$$\langle J\rangle = q(t_0^2 + (1/3)S(S+1))I(\mu) \quad (8)$$
$$-qt_2^2(B^2/T_{\rm eff})(1/6)S(S+1)\frac{{\rm d}I}{{\rm d}\mu}.$$

Current Noise— We now wish to compute the current-current correlator, $\langle J(\omega)J(-\omega)\rangle$. At second order in $\mathcal{H}_T$, we obtain the shot noise contribution. It is obtained by using Eq. (6) for J and computing the correlation function $\langle J(\omega)J(-\omega)\rangle$ using the Hamiltonian $\mathcal{H}_L + \mathcal{H}_R$, and no higher powers of $\mathcal{H}_T$. The result is $\langle J(\omega)J(-\omega)\rangle = 2q\langle J\rangle$, for $\omega, \omega_L << \mu$.

At fourth order in $\mathcal{H}_T$, many contributions are possible. The full calculation is quite involved, however the most important contribution is a "modulation" of the current. We consider a modulation of the current to be a term in $\langle J(\omega)J(-\omega)\rangle$ which is of order $\mu^{2(\alpha+1)}$. In contrast, for example, the shot noise is only of order $\mu^{\alpha+1}$.

Such modulation terms are obtained as follows [15]: we first consider S_z to be a classical field, and obtain the dependence of the current on S_z, giving $J = q(t_0^2 + t_2^2 S_z^2)I(\mu)$. Then, we consider how S_z varies in time, using the dynamics (4), to compute the correlation functions of S_z. The result is

$$\langle J(0)J(t)\rangle = q^2 t_2^4 I(\mu)^2 \langle S_z^2(0)S_z^2(t)\rangle. \quad (9)$$

For $S > 1/2$, this gives a modulation of the current at twice the Larmor frequency. However, there is no modulation at the Larmor frequency itself for any S.

Diagrammatically, we show a term giving such modulation in Fig. 1(a). Lines with arrows represent electron propagators, while pairs of straight lines without arrows represent spin propagators (here, simply to give some diagrammatic way of writing interactions with spins, we have chosen to use the standard represention of a spin operator as a pair of fermion operators). The curly lines the current operator, while the crosses represent the operator $\mathcal{H}_T$; since these two operators have straight lines leaving them, this indicates that we take the term $-qt_2T_0S_z$ in the current operator, and the term $t_2J_0S_z$ in $\mathcal{H}_T$. There are two pairs of lines connecting the bubbles, giving a modulation at twice the Larmor frequency. In Fig. 1(b), we show a similar diagram with only one pairs of lines connecting the bubbles; however, this diagram vanishes.

The fact that the modulation is not present at frequency ω_L will be true to all orders. In contrast, other terms in $\langle J(-\omega)J(\omega)\rangle$, which are lower order in μ and thus do not represent a modulation for this experimental setup, may show a peak at the Larmor frequency. These terms all depend on S_z having a time dependence on time scales of order μ^{-1} (the time over which a tunneling event occurs), and thus all of these terms will be suppressed by powers of B/μ and are therefore expected to be quite small, for this situation. In such terms, we would find diagrams with both current operators on the same electron bubble, rather than on different bubbles as was the case in Fig. 1.

However, for other situations, these terms may be important. Consider sequential tunneling from one lead to the other via a quantum dot [6], where the chemical potential in one lead is far higher in energy than the dot, and the chemical potential in the other lead is far lower in energy than the dot. In this case, the current through the system is largely independent of the exact difference, μ, between the chemical potentials in the two leads. Our considerations in this section relied on computing the order of terms in μ. However, in the given experimental setup with the quantum dot, the current is of order μ^0 and the time to tunnel from left to right lead is determined not by μ, but by the hopping matrix elements between the quantum dot and the leads. Then, it has been shown that [10] for tunneling through a quantum dot in this sequential tunneling regime, one can obtain a contribution to the current at the Larmor frequency which is comparable to the zero frequency component of the current. Thus, while in [6] terms were considered which violate time reversal symmetry, with slight modifications a signal at the Larmor frequency can be found using a Hamiltonian which respects time reversal symmetry.

More General Hamiltonians– The reason for a lack of modulation is that if the impurity spin is held fixed, with a static S_z, then the current depends only on S_z^2 and not on S_z itself. For the Hamiltonians considered above, this results from the fact that the current depends only on the modulus of the hopping coefficient

z above. However, this is true even in a more general case. Consider any Hamiltonian including potential, interaction, and position-dependent coupling of the electron current to S_z: $\mathcal{H} = \psi^\dagger(x)(\partial^2/2m + U(x))\psi(x) + \psi^\dagger(x)\psi(x)\psi^\dagger(y)\psi(y)V(x-y) + i\psi^\dagger(x)\{F(x), \partial_x\}\psi(x)S_z$. The spin-orbit coupling introduces an effective vector potential. If this vector potential has a non-vanishing curl, then it leads to a non-vanishing effective magnetic field and it *is* possible for the current to depend on the sign of S_z. However, in cases like those considered above, in which this effective vector potential acts only across a single tunnel junction and does not produce any magnetic field, the current does not depend on the sign of S_z, only on its magnitude. Further, in many cases with non-vanishing effective field, the current will still depend only on the sign of S_z, as must be determined on a case-by-case basis; for many systems, the magnetoresistance curve is symmetric in the magnetization, for example [17].

Example with Modulation— Let us finally give an example that does leads to a modulation at the Larmor frequency. Suppose the Hamiltonian is $\mathcal{H}_L+\mathcal{H}_R+\mathcal{H}_T+BS_x$, with $\mathcal{H}_T = t_0 T_0 + t_2^a J_0(S_j^a w_j) + t_2^b J_0(S_j^b w_j)$ and $\mathcal{H}_o = g^a B_i^a S_i^a + g^b B_i^b S_i^b$, where there are two impurity spins, a, b, with different g-factors, coupled to the current. Then, choosing w to lie along the z-direction, the current modulation is

$$\langle J(0)J(t)\rangle = I(\mu)^2 \langle \Omega(0)^2 \Omega(t)^2 \rangle, \tag{10}$$

where we define $\Omega = t_2^a S_z^a + t_2^b S_z^b$. Then, if the two spins have different precession frequencies $\omega_L^{1,2}$, the two spins will "beat" together and peaks will be observed at $\pm 2\omega_L^a, \pm 2\omega_L^b, \pm\omega_L^a \pm \omega_L^b$. If one spin precesses slowly, $\omega_L^b \approx 0$, then this gives a peak at ω_L^a. Alternately, the same result can be obtained by tunneling spin-polarized electrons, again giving rise to a modulation at ω_L.

Conclusion— In conclusion, we have shown that the assumption of time reversal symmetry severely constrains the ability to measure a spin via a tunneling current. For weak magnetic fields, where we can neglect possible Aharonov-Bohm phases, the modulation is observed at twice the Larmor frequency, rather than at the Larmor frequency. This result is fairly general for systems with a single tunnel junction; for systems with a quantum dot in the sequential tunneling regime, or other systems in which the tunneling current is largely independent of the voltage, a peak in the current-current correlator at the Larmor frequency is possible. While we have derived this result for a specific Hamiltonian (2), it holds for a more general Hamiltonian, if we take $\mathcal{H} = \mathcal{H}_L + \mathcal{H}_R + \mathcal{H}_T + BS_x$, with $\mathcal{H}_T$ given by Eq. (1).

Given that such a modulation is observed at the Larmor frequency, and assuming the Aharonov-Bohm phases are negligible, then either the system is in a sequential tunneling regime or else there are multiple spins present, enabling a beating of the precession frequencies of the two spins. However, in the latter case there should be a current modulation at twice the Larmor frequency, at least for impurity spins greater than 1/2, and thus an experimental measurement of that peak is suggested.

Acknowledgments— I thank D. Mozyrsky, I. Martin, and A. V. Balatsky for useful discussions. This work was supported by US DOE W-7405-ENG-36.

[1] M. Bode, M. Getzlaff, and R. Wiesendanger, Phys. Rev. Lett., **81**, 4256 (1998).
[2] Z. Nussinov, M. F. Crommie, and A. V. Balatsky, Phys. Rev. B **68**, 085402 (2003).
[3] M. Farle, Rep. Prog. Phys. **61**, 755 (1998).
[4] G. Prinz, Science **282**, 1660 (1998).
[5] D. Shachal and Y. Manassen, Phys. Rev. B **46**, 4795 (1992); Y. Manassen, I. Mukhupadyay, and N. Ramesh Rao, Phys. Rev. B **61** 16223 (2000).
[6] D. Mozyrsky et. al., Phys. Rev. B **66**, 161313 (2002).
[7] L. N. Bulaevski, M. Hruska, and G. Ortiz, Phys. Rev. B **68**, 125415 (2003).
[8] J.-X. Zhu and A. V. Balatsky, Phys. Rev. Lett, **89**, 286802 (2002).
[9] One possibility is a paramagnetic cluster near the tip of the tunneling system, for example. See A. V. Balatsky, Y. Manassen, and R. Salem, Phys. Rev. B **66**, 195416 (2002).
[10] D. Mozyrsky, private communication, to be published.
[11] L. D. Landau and E. M. Lifshitz, *Quantum Mechanics* (Butterworth-Heineman, Boston, 2000).
[12] V. F. Gantmakher and I. B. Levinson, *Carrier Scattering in Metals and Semiconductors* (Elesevier/Science, Amsterdam, 1987).
[13] A. Shnirman, D. Mozyrsky, and I. Martin, preprint cond-mat/0211618.
[14] C. L. Kane and M. P. A. Fisher, Phys. Rev. Lett. **68**, 1220 (1992).
[15] M. B. Hastings, I. Martin, and D. Mozyrsky, Phys. Rev. B **68**, 035101 (2003); D. Mozyrsky and I. Martin, Phys. Rev. Lett. **89**, 018301 (2002).
[16] L. V. Keldysh, Sov. Phys. JETP **20**, 1018 (1965).
[17] Effects of different relaxation mechanisms and other possible spin-orbit coupling terms such as Rashba terms will be discussed elsewhere; M. B. Hastings, I. Martin, and D. Mozyrsky, to be published.



FIG. 1. (a) Example diagram showing modulation at $2\omega_L$. (b) Diagram which vanishes.